

## DIAGNOSTIC IMAGING IN HEMATOLOGY

### **Rapid and Severe Covid-19 Pneumonia with Severe Acute Chest Syndrome in a Sick Cell Patient Successfully Treated with Tocilizumab**

Gonzalo De Luna<sup>1+</sup> M.D.; Anoosha Habibi<sup>1+</sup> M.D. Ph.D.; Jean François Deux<sup>2</sup> M.D. Ph.D.; Martin Colard<sup>1</sup> M.D. Ph.D.; Anne Laure Pham Hung d'Alexandry d'Orengiani<sup>3</sup> Ph.D.; Frédéric Schlemmer<sup>4</sup> M.D., Ph.D.; Nizar Joher<sup>5</sup> M.D.; Christian Kassasseya<sup>6</sup> M.D.; Jean Michel Pawlotsky<sup>7</sup> M.D. Ph.D.; Clément Ourghanlian<sup>8</sup> M.D., Marc Michel<sup>9</sup> M.D. Ph.D.; Armand Mekontso-Dessap<sup>10</sup> M.D. Ph.D.; Pablo Bartolucci<sup>1\*</sup> M.D. Ph.D..

<sup>1</sup> Sick Cell Referral Center, Department of Internal Medicine, Henri-Mondor University Hospital- UPEC, AP-HP, Créteil, France

<sup>2</sup> Department of Radiology, Henri Mondor Hospital/AP-HP, F-94010, Créteil, France

<sup>3</sup> MCGRE, APHP, CHU Henri Mondor, Créteil, France

<sup>4</sup> Service de Pneumologie, DHU A-TVB, APHP, CHU Henri Mondor, Créteil, France

<sup>5</sup> Nephrology and Renal Transplantation Department, Institut Francilien de Recherche en Néphrologie et Transplantation (IFRNT), Groupe Hospitalier Henri-Mondor/Albert-Chenevier, Créteil, France

<sup>6</sup> Emergency Department, APHP, CHU Henri Mondor, Créteil, France

<sup>7</sup> National Reference Center for Viral Hepatitis B, C and D, Department of Virology, Hôpital Henri Mondor, Université Paris-Est, Créteil, France

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ajh.25833

<sup>8</sup> AP-HP, Hôpital Henri Mondor, Pharmacie, F-94000, Créteil, France

<sup>9</sup> Department of Internal Medicine, National Referral Center for Adult Immune Cytopenias, Henri Mondor University Hospital, Assistance Publique Hôpitaux de Paris, Université Paris-Est Créteil, France

<sup>10</sup> Department of Intensive Care, Henri Mondor University Hospital, Assistance Publique Hôpitaux de Paris, Université Paris-Est Créteil, France

<sup>+</sup> These authors equally contributed to this manuscript

<sup>\*</sup> Corresponding author:

Pablo Bartolucci; Phone: + (00)33.1.49.81.24.43; E-mail address: [pablo.bartolucci@aphp.fr](mailto:pablo.bartolucci@aphp.fr)

KEYWORDS: Covid-19; sickle cell disease; acute chest syndrome; anti-human interleukin-6 receptor monoclonal antibody

Accepted Article

On March 25, 2020, presence of multifocal vaso-occlusive crises (VOC) for the past 24 hours was determined in a 45-year-old male patient with homozygous sickle cell disease (SCD) by the "DREPADOM" network, a phone consultation clinical monitoring and ambulatory care of SCD patients set up in our SCD referral center since the outbreak of the Covid-19 epidemic in France. Prior medical history included sickle cell nephropathy with tubular acidosis, ischemic retinopathy, priapism and cardiac remodeling. Past History for VOC or Acute Chest Syndrome (ACS) was negative for the past 10 years, except for a brief hospitalization in February 2020 for sub-segmental pulmonary embolism, secondary to ACS and treated with Rivaroxaban. Hydroxyurea treatment was scheduled after sperm cryo-preservation but had not yet started at the time of the Covid-19 hospitalization. After the DREPADOM screening, he was admitted for multifocal VOC with normal pulmonary findings, no dyspnea, no diarrhea, no anosmia, no cough, no fever and oxygen saturation (SpO<sub>2</sub>) at 98%. On day 1 of hospitalization, the patient developed fever (38.5°C) and SpO<sub>2</sub> dropped to 91% with crackles at pulmonary auscultation. Antibiotic therapy was immediately started with Amoxicillin-clavulanic acid and the patient received supplemental oxygen through a nasal cannula at a rate of 3 L/min. Hydroxychloroquine treatment at a dosage of 200mg orally every 8 hours was instituted, while results of the real-time reverse-transcription–polymerase-chain-reaction (RT-PCR) assay were still pending. The electrocardiogram showed a QT interval at 390ms. On day 2, the patient's general condition rapidly deteriorated and SpO<sub>2</sub> dropped to 80%. Supplemental oxygen through venturi mask at a rate of 15L/min and a 100% fraction of inspired oxygen maintained the SpO<sub>2</sub> at 91%. Surprisingly, the patient presented a respiratory rate of 19 breaths/min. Notable laboratory values were: Hemoglobin 7 g/L, reticulocytes 186 Giga/L, leucocytes 20 Giga/L, C-reactive protein at 189 mg/L, serum ferritin 3271 µg/L and creatinine clearance (DFG CKD EPI) 120 mL/min/1.73m<sup>2</sup>. Computerized tomography (CT) of the chest displayed abnormalities, consistent with a Covid-19-induced pneumonia and ACS.

**(Figure 1).** The RT-PCR assay for the Covid-19 diagnosis was positive. Treatment with one pulse of intravenous Tocilizumab at a dosage of 8 mg/kg was administered. On day 3, a clear improvement of the patient's general condition was observed, with a SpO<sub>2</sub> at 97% by supplemental oxygen through a nasal cannula at a rate of 3 L/min and no fever. On day 4, blood transfusion was performed due to the ACS condition. On day 5, the patient was discharged and referred back for ambulatory care to the DREPADOM structure.

SCD is a serious genetic condition that shortens life expectancy. It affects more than 30,000 people in France, 50% of whom are located in the Ile de France region(2). A severe complication of SCD is ACS, that can be triggered by infectious complications(1). The Influenza H1N1 epidemic had a 17% rate of hospitalization in intensive care units for the SCD population(3,4). Covid-19 and the associated Acute Respiratory Distress Syndrome (ARDS), represent a significant mortality risk for SCD patients. Extracorporeal oxygenation (ECMO), which is often required in ARDS, is associated in SCD patients with catastrophic prognosis (70% mortality rate)(5). IL-6 is a multifunctional cytokine that plays a central role in host defense mechanisms. Abnormally high plasma values of IL-6 have been reported in SCD patients at steady (healthy) state(6). IL-6 and C reactive protein are elevated during VOC. Inflammation contributes to the sickle red blood cells adhesion process involved in vaso-occlusive pathophysiology(8). The SARS-CoV S protein induces direct up-regulation of IL-6(7), IL-1 and TNF $\alpha$ , some of the most potent pro-inflammatory cytokines. Tocilizumab (TCZ) is an anti-human IL-6 receptor monoclonal antibody that inhibits signal transduction by binding sIL-6R and mIL-6R. Despite the lack of clinical trials on TCZ efficacy and safety for Covid-19 treatment, it was recently approved in China for patients affected by severe Covid-19 pulmonary complications. Preliminary data from an observational study conducted in China on 21 severe cases receiving TCZ(9), showed improvement of clinical and radiological outcomes. Early antiviral strategies at the onset of the infection should be

considered for high risk patients. For critically ill patients, therapy directed toward the chemokine release syndrome is required. For our patient, given the prior history of severe SCD and the potential risks, treatment with hydroxychloroquine and TCZ were initiated, with a positive resolution. More studies are needed to determine the proper therapy for COVID-19 in patients affected by SCD.

#### FUNDING

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

#### COMPETING INTERESTS

Pr BARTOLUCCI reports being a consultant for F. Hoffmann-La Roche. No other potential conflict of interest relevant to this letter was reported.

#### FIGURE LEGEND

**Figure 1.** CT scan of the chest: Acute chest syndrome and Covid-19-induced pneumonia.

Axial image of chest obtained with a soft-tissue windows at the level of the lower lobes (panel A) evidenced areas of consolidation located at the posterior part of the lung (arrows). An axial image with the same windowing obtained at the upper part of the lungs (panel B) showed a right small pleural effusion in the upper part of the great pleural cavity (arrow). Axial image located at the same level than panel A with lung windows evidenced areas of ground-glass opacities (Panel C, arrows) in the lower lungs with regards to areas of consolidation, but also in the middle lobe (Panel C, arrowhead) and in the upper right lobe (Panel D). Coronal reconstruction (Panel E) confirmed areas of ground-glass opacities (arrow) and areas of consolidation with air bronchograms (arrowhead). Magnification of a CT image with lung windows acquired at the middle part of the lungs showing a

crazy-paving pattern with ground-glass opacities and interlobar septal thickening (Panel F; arrowhead).

## REFERENCES:

1. Vichinsky EP, Neumayr LD, Earles AN, Williams R, Lennette ET, Dean D, et al. Causes and outcomes of the acute chest syndrome in sickle cell disease. National Acute Chest Syndrome Study Group. *N Engl J Med*. 2000 Jun 22;342(25):1855–65.
2. Article - Bulletin épidémiologique hebdomadaire [Internet]. [cited 2020 Apr 3]. Available from: [http://beh.santepubliquefrance.fr/beh/2015/8/2015\\_8\\_2.html](http://beh.santepubliquefrance.fr/beh/2015/8/2015_8_2.html)
3. Bundy DG, Strouse JJ, Casella JF, Miller MR. Burden of influenza-related hospitalizations among children with sickle cell disease. *Pediatrics*. 2010 Feb;125(2):234–43.
4. Strouse JJ, Reller ME, Bundy DG, Amoako M, Cancio M, Han RN, et al. Severe pandemic H1N1 and seasonal influenza in children and young adults with sickle cell disease. *Blood*. 2010 Nov 4;116(18):3431–4.
5. Boissier F, Bagate F, Schmidt M, Labbé V, Kimmoun A, Fartoukh M, et al. Extracorporeal Life Support for Severe Acute Chest Syndrome in Adult Sickle Cell Disease: A Preliminary Report. *Crit Care Med*. 2019;47(3):e263–5.
6. Taylor SC, Shacks SJ, Mitchell RA, Banks A. Serum interleukin-6 levels in the steady state of sickle cell disease. *J Interferon Cytokine Res Off J Int Soc Interferon Cytokine Res*. 1995 Dec;15(12):1061–4.
7. Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet Lond Engl*. 2020 28;395(10229):1033–4.
8. Conran N, Belcher JD. Inflammation in sickle cell disease. *Clin Hemorheol Microcirc*. 2018;68(2–3):263–99.
9. Effective treatment of severe COVID-19 patients with Tocilizumab - La SFAR [Internet]. Société Française d'Anesthésie et de Réanimation. 2020 [cited 2020 Apr 3]. Available from: <https://sfar.org/effective-treatment-of-severe-covid-19-patients-with-tocilizumab/>



